

# Double Field Theory

CMH & Barton Zwiebach

arXiv:0904.4664,  
0908.1792

CMH & BZ & Olaf Hohm

arXiv:1003.5027, 1006.4664

# Double Field Theory


- From sector of String Field Theory, features stringy physics, including T-duality
- Strings see a doubled space-time
- Doubled space fully dynamic
- Interesting geometry
- Relations to generalised geometry which doubles tangent space, but not space itself.

# T-duality

- Takes  $S^1$  of radius  $R$  to  $S^1$  of radius  $1/R$
- Exchanges momentum  $p$  and winding  $w$
- Exchanges  $S^1$  coordinate  $X$  and dual  $S^1$  coordinate  $\tilde{X}$
- Acts on “doubled circle” with coordinates  $(X, \tilde{X})$
- On  $d$  torus, T-duality group  $O(d, d; \mathbb{Z})$
- Automorphisms of conformal field theory

# T-Duality

- Space has d-torus fibration
- G,B on fibres
- T-Duality  $O(d,d;\mathbb{Z})$ , mixes G,B
- Mixes Momentum and Winding
- Changes geometry and topology



$$E \rightarrow (aE + b)(cE + d)^{-1}$$

$$h = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in O(d, d; \mathbb{Z}) \quad E_{ij} = G_{ij} + B_{ij}$$

On circle, radius R:  $O(1, 1; \mathbb{Z}) = \mathbb{Z}_2 : R \mapsto \frac{1}{R}$

# Strings on $T^d$

$$X = X_L(\sigma + \tau) + X_R(\sigma - \tau), \quad \tilde{X} = X_L - X_R$$

$X$  conjugate to momentum,  $\tilde{X}$  to winding no.

$$dX = *d\tilde{X} \quad \partial_a X = \epsilon_{ab} \partial^b \tilde{X}$$

# Strings on $T^d$

$$X = X_L(\sigma + \tau) + X_R(\sigma - \tau), \quad \tilde{X} = X_L - X_R$$

$X$  conjugate to momentum,  $\tilde{X}$  to winding no.

$$dX = *d\tilde{X} \quad \partial_a X = \epsilon_{ab} \partial^b \tilde{X}$$

Need “auxiliary”  $\tilde{X}$  for interacting theory

i) Vertex operators  $e^{ik_L \cdot X_L}, e^{ik_R \cdot X_R}$

ii) String field **Kugo & Zwiebach**  $\Phi[x, \tilde{x}, a, \tilde{a}]$

# Strings on $T^d$

$$X = X_L(\sigma + \tau) + X_R(\sigma - \tau), \quad \tilde{X} = X_L - X_R$$

$X$  conjugate to momentum,  $\tilde{X}$  to winding no.

$$dX = *d\tilde{X} \quad \partial_a X = \epsilon_{ab} \partial^b \tilde{X}$$

Need “auxiliary”  $\tilde{X}$  for interacting theory

i) Vertex operators  $e^{ik_L \cdot X_L}, e^{ik_R \cdot X_R}$

ii) String field **Kugo & Zwiebach**  $\Phi[x, \tilde{x}, a, \tilde{a}]$

**Doubled Torus** 2d coordinates

Transform linearly under  $O(d, d; \mathbb{Z})$

$$X \equiv \begin{pmatrix} \tilde{x}^i \\ x^i \end{pmatrix}$$

Doubled sigma model **CMH 0406102**

# Strings on a Torus



- States: momentum  $p$ , winding  $w$
- String: Infinite set of fields  $\psi(p, w)$
- Fourier transform to doubled space:  $\psi(x, \tilde{x})$
- “Double Field Theory” from closed string field theory. Some non-locality in doubled space
- Subsector? e.g.  $g_{ij}(x, \tilde{x})$ ,  $b_{ij}(x, \tilde{x})$ ,  $\phi(x, \tilde{x})$



# Double Field Theory

- Double field theory on doubled torus
- General solution of string theory: involves doubled fields  $\psi(x, \tilde{x})$
- DFT needed for non-geometric backgrounds
- *Real* dependence on *full* doubled geometry, dual dimensions not auxiliary or gauge artifact. Double geom. *physical* and *dynamical*

- Novel symmetry, reduces to diffeos + B-field trans. in *any* half-dimensional subtorus
- Backgrounds depending on  $\{x^a\}$  seen by particles, on  $\{\tilde{x}_a\}$  seen by winding modes.
- Captures exotic and complicated structure of interacting string: **Non-polynomial, algebraic structure homotopy Lie algebra, cocycles.**
- T-duality symmetry manifest
- Formalism for T-folds etc
- Generalised T-duality: no isometries needed

**Earlier work on double fields:  
Siegel, Tseytlin**

# Strings on Torus

$$D=n+d$$

Target space

$$\mathbb{R}^{n-1,1} \times T^d$$

Coordinates

$$x^i = (x^\mu, x^a)$$

Momenta

$$p_i = (p_\mu, p_a)$$

Winding

$$w^i = (w^\mu, w^a)$$

Dual coordinates (conjugate to winding)

$$\tilde{x}_i = (\tilde{x}_\mu, \tilde{x}_a)$$

Constant metric and B-field

$$E_{ij} = G_{ij} + B_{ij}$$

## Compact dimensions

$p_a, w^a$  discrete, in Narain lattice,  $x^a, \tilde{x}_a$  periodic

Non-compact dimensions  $x^\mu, p_\mu$  continuous

Usually take  $w^\mu = 0$  so  $\frac{\partial}{\partial \tilde{x}_\mu} = 0$ , fields  $\psi(x^\mu, x^a, \tilde{x}_a)$

# T-Duality

- Interchanges momentum and winding
- Equivalence of string theories on dual backgrounds with very different geometries
- String field theory symmetry, provided fields depend on both  $x, \tilde{x}$  **Kugo, Zwiebach**
- For fields  $\psi(x^\mu)$  not  $\psi(x^\mu, x^a, \tilde{x}_a)$  **Buscher**
- Generalise to fields  $\psi(x^\mu, x^a, \tilde{x}_a)$

**Generalised T-duality**

**Dabholkar & CMH**

# String Field Theory on Minkowski Space

Closed SFT:  
Zwiebach

String field  $\Phi[X(\sigma), c(\sigma)]$

$X^i(\sigma) \rightarrow x^i$ , oscillators

Expand to get infinite set of fields

$g_{ij}(x), b_{ij}(x), \phi(x), \dots, C_{ijk\dots l}(x), \dots$

Integrating out massive fields gives field theory for

$g_{ij}(x), b_{ij}(x), \phi(x)$

# String Field Theory on a torus

String field  $\Phi[X(\sigma), c(\sigma)]$

$X^i(\sigma) \rightarrow x^i, \tilde{x}_i$ , oscillators

Expand to get infinite set of double fields

$g_{ij}(x, \tilde{x}), b_{ij}(x, \tilde{x}), \phi(x, \tilde{x}), \dots, C_{ijk\dots l}(x, \tilde{x}), \dots$

Seek **double field theory** for

$g_{ij}(x, \tilde{x}), b_{ij}(x, \tilde{x}), \phi(x, \tilde{x})$

# Free Field Equations (B=0)

$$L_0 + \bar{L}_0 = 2$$

$$p^2 + w^2 = N + \bar{N} - 2$$

$$L_0 - \bar{L}_0 = 0$$

$$p_i w^i = N - \bar{N}$$

# Free Field Equations (B=0)

$$L_0 + \bar{L}_0 = 2$$

$$p^2 + w^2 = N + \bar{N} - 2$$

Treat as field equation, kinetic operator in doubled space

$$G^{ij} \frac{\partial^2}{\partial x^i \partial x^j} + G_{ij} \frac{\partial^2}{\partial \tilde{x}_i \partial \tilde{x}_j}$$

$$L_0 - \bar{L}_0 = 0$$

$$p_i w^i = N - \bar{N}$$

Treat as constraint on double fields

$$\Delta \equiv \frac{\partial^2}{\partial x^i \partial \tilde{x}_i} \quad (\Delta - \mu)\psi = 0$$



# Free Field Equations (B=0)

$$L_0 + \bar{L}_0 = 2$$

$$p^2 + w^2 = N + \bar{N} - 2$$

Treat as field equation, kinetic operator in doubled space

$$G^{ij} \frac{\partial^2}{\partial x^i \partial x^j} + G_{ij} \frac{\partial^2}{\partial \tilde{x}_i \partial \tilde{x}_j}$$

Laplacian for metric

$$L_0 - \bar{L}_0 = 0$$

$$p_i w^i = N - \bar{N}$$

$$ds^2 = G_{ij} dx^i dx^j + G^{ij} d\tilde{x}_i d\tilde{x}_j$$

Treat as constraint on double fields

Laplacian for metric

$$\Delta \equiv \frac{\partial^2}{\partial x^i \partial \tilde{x}_i} \quad (\Delta - \mu)\psi = 0$$

$$ds^2 = dx^i d\tilde{x}_i$$

$$g_{ij}(x, \tilde{x}), b_{ij}(x, \tilde{x}), \phi(x, \tilde{x})$$

$$N = \bar{N} = 1$$

$$p^2 + w^2 = 0$$

$$p \cdot w = 0$$

**“Double Massless”**

Constrained fields  $\psi(x^\mu, x^a, \tilde{x}_a)$

$$(\Delta - \mu)\psi = 0$$

**Momentum space**  $\psi(p_\mu, p_a, w^a)$   $\Delta = p_a w^a$

**Momentum space: Dimension  $n+2d$**

**Cone:  $p_a w^a = 0$  or hyperboloid:  $p_a w^a = \mu$**

**dimension  $n+2d-1$**

**DFT: fields on cone or hyperboloid, with discrete  $p, w$**

**Problem: naive product of fields on cone do not lie on cone. Vertices need projectors**

Restricted fields: Fields that depend on  $d$  of  $2d$  torus

**momenta, e.g.  $\psi(p_\mu, p_a)$  or  $\psi(p_\mu, w^a)$**

**Simple subsector, no projectors needed, no cocycles.**

# Torus Backgrounds

$$G_{ij} = \begin{pmatrix} \eta_{\mu\nu} & 0 \\ 0 & G_{ab} \end{pmatrix}, \quad B_{ij} = \begin{pmatrix} 0 & 0 \\ 0 & B_{ab} \end{pmatrix} \quad E_{ij} \equiv G_{ij} + B_{ij}$$

Fluctuations  $e_{ij} = h_{ij} + b_{ij}$

Take  $B_{ij} = 0$   $\tilde{\partial}_i \equiv G_{ik} \frac{\partial}{\partial \tilde{x}_k}$

# Torus Backgrounds

$$G_{ij} = \begin{pmatrix} \eta_{\mu\nu} & 0 \\ 0 & G_{ab} \end{pmatrix}, \quad B_{ij} = \begin{pmatrix} 0 & 0 \\ 0 & B_{ab} \end{pmatrix} \quad E_{ij} \equiv G_{ij} + B_{ij}$$

Fluctuations  $e_{ij} = h_{ij} + b_{ij}$

Take  $B_{ij} = 0$   $\tilde{\partial}_i \equiv G_{ik} \frac{\partial}{\partial \tilde{x}_k}$

Usual action  $\int dx \sqrt{-g} e^{-2\phi} \left[ R + 4(\partial\phi)^2 - \frac{1}{12} H^2 \right]$

Quadratic part  $\int dx L[h, b, d; \partial]$

$$e^{-2d} = e^{-2\phi} \sqrt{-g}$$

(d invariant under usual T-duality)

# Double Field Theory Action

$$S^{(2)} = \int [dx d\tilde{x}] \left[ L[h, b, d; \partial] + L[-h, -b, d; \tilde{\partial}] \right. \\ \left. + (\partial_k h^{ik})(\tilde{\partial}^j b_{ij}) + (\tilde{\partial}^k h_{ik})(\partial_j b^{ij}) - 4d \partial^i \tilde{\partial}^j b_{ij} \right]$$

**Action + dual action + strange mixing terms**

# Double Field Theory Action

$$S^{(2)} = \int [dx d\tilde{x}] \left[ L[h, b, d; \partial] + L[-h, -b, d; \tilde{\partial}] \right. \\ \left. + (\partial_k h^{ik})(\tilde{\partial}^j b_{ij}) + (\tilde{\partial}^k h_{ik})(\partial_j b^{ij}) - 4d \partial^i \tilde{\partial}^j b_{ij} \right]$$

Action + dual action + strange mixing terms

$$\delta h_{ij} = \partial_i \epsilon_j + \partial_j \epsilon_i + \tilde{\partial}_i \tilde{\epsilon}_j + \tilde{\partial}_j \tilde{\epsilon}_i ,$$

$$\delta b_{ij} = -(\tilde{\partial}_i \epsilon_j - \tilde{\partial}_j \epsilon_i) - (\partial_i \tilde{\epsilon}_j - \partial_j \tilde{\epsilon}_i) ,$$

$$\delta d = -\partial \cdot \epsilon + \tilde{\partial} \cdot \tilde{\epsilon} . \quad \text{Invariance needs constraint}$$

Diffeos and B-field transformations mixed.

Invariant cubic action found for full DFT of (h,b,d)

# T-Duality Transformations of Background

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in O(d, d; \mathbb{Z}) \quad \text{T-duality}$$

$$E' = (aE + b)(cE + d)^{-1}$$

$$X \equiv \begin{pmatrix} \tilde{x}_i \\ x^i \end{pmatrix} \quad \text{transforms as a vector}$$

$$X' = \begin{pmatrix} \tilde{x}' \\ x' \end{pmatrix} = gX = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \tilde{x} \\ x \end{pmatrix}$$



# T-Duality is a Symmetry of the Action

Fields  $e_{ij}(x, \tilde{x}), d(x, \tilde{x})$

Background  $E_{ij}$

$$E' = (aE + b)(cE + d)^{-1}$$

$$X' = \begin{pmatrix} \tilde{x}' \\ x' \end{pmatrix} = gX = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \tilde{x} \\ x \end{pmatrix}$$

Action invariant if:

$$e_{ij}(X) = M_i^k \bar{M}_j^l e'_{kl}(X')$$

$$M \equiv d^t - E c^t$$

$$d(X) = d'(X')$$

$$\bar{M} \equiv d^t + E^t c^t$$

**With general momentum and winding dependence!**

# Projectors and Cocycles

Naive product of constrained fields doesn't satisfy constraint

$$L_0^- \Psi_1 = 0, L_0^- \Psi_2 = 0 \quad \text{but} \quad L_0^- (\Psi_1 \Psi_2) \neq 0$$

$$\Delta A = 0, \Delta B = 0 \quad \text{but} \quad \Delta(AB) \neq 0$$

String product has explicit projection

Leads to a symmetry that is not a Lie algebra, but is a homotopy lie algebra.

# Projectors and Cocycles

Naive product of constrained fields doesn't satisfy constraint

$$L_0^- \Psi_1 = 0, L_0^- \Psi_2 = 0 \quad \text{but} \quad L_0^- (\Psi_1 \Psi_2) \neq 0$$

$$\Delta A = 0, \Delta B = 0 \quad \text{but} \quad \Delta(AB) \neq 0$$

String product has explicit projection

Leads to a symmetry that is not a Lie algebra, but is a homotopy lie algebra.

Double field theory requires projections.

SFT has non-local cocycles in vertices, DFT should too  
Cocycles and projectors not needed in cubic action

General double fields

$$\psi(x, \tilde{x})$$

Fields on Spacetime M

$$\psi(x)$$

Restricted Fields on N, T-dual to M

$$\psi(x')$$

M, N null wrt  $O(D, D)$  metric  $ds^2 = dx^i d\tilde{x}_i$

**Subsector with fields and parameters all restricted to M or N**

- Constraint satisfied on all fields and products of fields
- No projectors or cocycles
- T-duality covariant: independent of choice of N
- Can find full non-linear form of gauge transformations
- Full gauge algebra, full non-linear action

# Restricted DFT

Double fields restricted to null D-dimensional subspace N  
T-duality “rotates” N to N’

Background independent fields: g,b,d:

$$\mathcal{E} \equiv E + \left(1 - \frac{1}{2} e\right)^{-1} e \quad \mathcal{E}_{ij} = g_{ij} + b_{ij}$$

## **O(D,D) Covariant Notation**

$$X^M \equiv \begin{pmatrix} \tilde{x}_i \\ x^i \end{pmatrix} \quad \partial_M \equiv \begin{pmatrix} \partial^i \\ \partial_i \end{pmatrix}$$
$$\eta_{MN} = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \quad M = 1, \dots, 2D$$

# Generalised T-duality transformations:

$$X'^M \equiv \begin{pmatrix} \tilde{x}'_i \\ x'^i \end{pmatrix} = h X^M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \tilde{x}_i \\ x^i \end{pmatrix}$$

$h$  in  $O(d,d;\mathbf{Z})$  acts on toroidal coordinates only

$$\mathcal{E}'(X') = (a\mathcal{E}(X) + b)(c\mathcal{E}(X) + d)^{-1}$$

$$d'(X') = d(X)$$

Buscher if fields independent of toroidal coordinates  
Generalisation to case without isometries

# O(D,D)

Non-compact dimensions  $x^\mu, p_\mu$  continuous

Strings: take  $w^\mu = 0$  so  $\frac{\partial}{\partial \tilde{x}_\mu} = 0$ , fields  $\psi(x^\mu, x^a, \tilde{x}_a)$

For DFT, if we allow dependence on  $\tilde{x}_\mu$

DFT invariant under

$$O(n, n) \times O(d, d; \mathbb{Z})$$

Subgroup of O(D,D) preserving periodicities

O(D,D) is symmetry if all directions non-compact:  
theory has formal O(D,D) covariance

# Generalised Metric Formulation

$$\mathcal{H}_{MN} = \begin{pmatrix} g^{ij} & -g^{ik}b_{kj} \\ b_{ik}g^{kj} & g_{ij} - b_{ik}g^{kl}b_{lj} \end{pmatrix} .$$

2 Metrics on double space

$\mathcal{H}_{MN}, \eta_{MN}$



# Generalised Metric Formulation

$$\mathcal{H}_{MN} = \begin{pmatrix} g^{ij} & -g^{ik}b_{kj} \\ b_{ik}g^{kj} & g_{ij} - b_{ik}g^{kl}b_{lj} \end{pmatrix}.$$

2 Metrics on double space  $\mathcal{H}_{MN}, \eta_{MN}$

$$\mathcal{H}^{MN} \equiv \eta^{MP}\mathcal{H}_{PQ}\eta^{QN}$$

Constrained metric  $\mathcal{H}^{MP}\mathcal{H}_{PN} = \delta^M_N$

# Generalised Metric Formulation

$$\mathcal{H}_{MN} = \begin{pmatrix} g^{ij} & -g^{ik}b_{kj} \\ b_{ik}g^{kj} & g_{ij} - b_{ik}g^{kl}b_{lj} \end{pmatrix}.$$

2 Metrics on double space

$$\mathcal{H}_{MN}, \eta_{MN}$$

$$\mathcal{H}^{MN} \equiv \eta^{MP}\mathcal{H}_{PQ}\eta^{QN}$$

Constrained metric

$$\mathcal{H}^{MP}\mathcal{H}_{PN} = \delta^M_N$$

Covariant Transformation

$$h^P_M h^Q_N \mathcal{H}'_{PQ}(X') = \mathcal{H}_{MN}(X)$$

$$X' = hX \quad h \in O(D, D)$$

## O(D,D) covariant action

$$S = \int dx d\tilde{x} e^{-2d} L$$

$$L = \frac{1}{8} \mathcal{H}^{MN} \partial_M \mathcal{H}^{KL} \partial_N \mathcal{H}_{KL} - \frac{1}{2} \mathcal{H}^{MN} \partial_N \mathcal{H}^{KL} \partial_L \mathcal{H}_{MK} \\ - 2 \partial_M d \partial_N \mathcal{H}^{MN} + 4 \mathcal{H}^{MN} \partial_M d \partial_N d$$

## O(D,D) covariant action

$$S = \int dx d\tilde{x} e^{-2d} L$$

$$L = \frac{1}{8} \mathcal{H}^{MN} \partial_M \mathcal{H}^{KL} \partial_N \mathcal{H}_{KL} - \frac{1}{2} \mathcal{H}^{MN} \partial_N \mathcal{H}^{KL} \partial_L \mathcal{H}_{MK} \\ - 2 \partial_M d \partial_N \mathcal{H}^{MN} + 4 \mathcal{H}^{MN} \partial_M d \partial_N d$$

**L cubic! Indices raised and lowered with  $\eta$**

## O(D,D) covariant action

$$S = \int dx d\tilde{x} e^{-2d} L$$

$$L = \frac{1}{8} \mathcal{H}^{MN} \partial_M \mathcal{H}^{KL} \partial_N \mathcal{H}_{KL} - \frac{1}{2} \mathcal{H}^{MN} \partial_N \mathcal{H}^{KL} \partial_L \mathcal{H}_{MK} \\ - 2 \partial_M d \partial_N \mathcal{H}^{MN} + 4 \mathcal{H}^{MN} \partial_M d \partial_N d$$

## Gauge Transformation

$$\delta_\xi \mathcal{H}^{MN} = \xi^P \partial_P \mathcal{H}^{MN} \\ + (\partial^M \xi_P - \partial_P \xi^M) \mathcal{H}^{PN} + (\partial^N \xi_P - \partial_P \xi^N) \mathcal{H}^{MP}$$

## O(D,D) covariant action

$$S = \int dx d\tilde{x} e^{-2d} L$$

$$L = \frac{1}{8} \mathcal{H}^{MN} \partial_M \mathcal{H}^{KL} \partial_N \mathcal{H}_{KL} - \frac{1}{2} \mathcal{H}^{MN} \partial_N \mathcal{H}^{KL} \partial_L \mathcal{H}_{MK} \\ - 2 \partial_M d \partial_N \mathcal{H}^{MN} + 4 \mathcal{H}^{MN} \partial_M d \partial_N d$$

## Gauge Transformation

$$\delta_\xi \mathcal{H}^{MN} = \xi^P \partial_P \mathcal{H}^{MN} \\ + (\partial^M \xi_P - \partial_P \xi^M) \mathcal{H}^{PN} + (\partial^N \xi_P - \partial_P \xi^N) \mathcal{H}^{MP}$$

Rewrite as “Generalised Lie Derivative”

$$\delta_\xi \mathcal{H}^{MN} = \hat{\mathcal{L}}_\xi \mathcal{H}^{MN}$$

# Generalised Lie Derivative

$$A_{N_1 \dots}^{M_1 \dots}$$

$$\begin{aligned} \hat{\mathcal{L}}_{\xi} A_M^N &\equiv \xi^P \partial_P A_M^N \\ &+ (\partial_M \xi^P - \partial^P \xi_M) A_P^N + (\partial^N \xi_P - \partial_P \xi^N) A_M^P \end{aligned}$$

# Generalised Lie Derivative

$$A_{N_1 \dots}^{M_1 \dots}$$

$$\begin{aligned} \widehat{\mathcal{L}}_{\xi} A_M^N &\equiv \xi^P \partial_P A_M^N \\ &+ (\partial_M \xi^P - \partial^P \xi_M) A_P^N + (\partial^N \xi_P - \partial_P \xi^N) A_M^P \end{aligned}$$

$$\begin{aligned} \widehat{\mathcal{L}}_{\xi} A_M^N &= \mathcal{L}_{\xi} A_M^N - \eta^{PQ} \eta_{MR} \partial_Q \xi^R A_P^N \\ &+ \eta_{PQ} \eta^{NR} \partial_R \xi^Q A_M^P \end{aligned}$$



# Generalized scalar curvature

$$\begin{aligned}\mathcal{R} \equiv & 4\mathcal{H}^{MN}\partial_M\partial_N d - \partial_M\partial_N\mathcal{H}^{MN} \\ & - 4\mathcal{H}^{MN}\partial_M d\partial_N d + 4\partial_M\mathcal{H}^{MN}\partial_N d \\ & + \frac{1}{8}\mathcal{H}^{MN}\partial_M\mathcal{H}^{KL}\partial_N\mathcal{H}_{KL} - \frac{1}{2}\mathcal{H}^{MN}\partial_M\mathcal{H}^{KL}\partial_K\mathcal{H}_{NL}\end{aligned}$$

# Generalized scalar curvature

$$\begin{aligned}\mathcal{R} \equiv & 4\mathcal{H}^{MN}\partial_M\partial_N d - \partial_M\partial_N\mathcal{H}^{MN} \\ & - 4\mathcal{H}^{MN}\partial_M d\partial_N d + 4\partial_M\mathcal{H}^{MN}\partial_N d \\ & + \frac{1}{8}\mathcal{H}^{MN}\partial_M\mathcal{H}^{KL}\partial_N\mathcal{H}_{KL} - \frac{1}{2}\mathcal{H}^{MN}\partial_M\mathcal{H}^{KL}\partial_K\mathcal{H}_{NL}\end{aligned}$$

$$S = \int dx d\tilde{x} e^{-2d} \mathcal{R}$$

# Generalized scalar curvature

$$\begin{aligned}\mathcal{R} \equiv & 4\mathcal{H}^{MN}\partial_M\partial_N d - \partial_M\partial_N\mathcal{H}^{MN} \\ & - 4\mathcal{H}^{MN}\partial_M d\partial_N d + 4\partial_M\mathcal{H}^{MN}\partial_N d \\ & + \frac{1}{8}\mathcal{H}^{MN}\partial_M\mathcal{H}^{KL}\partial_N\mathcal{H}_{KL} - \frac{1}{2}\mathcal{H}^{MN}\partial_M\mathcal{H}^{KL}\partial_K\mathcal{H}_{NL}\end{aligned}$$

$$S = \int dx d\tilde{x} e^{-2d} \mathcal{R}$$

Gauge Symmetry

$$\delta_\xi \mathcal{R} = \hat{\mathcal{L}}_\xi \mathcal{R} = \xi^M \partial_M \mathcal{R}$$

$$\delta_\xi e^{-2d} = \partial_M (\xi^M e^{-2d})$$

# Generalized scalar curvature

$$\begin{aligned}\mathcal{R} \equiv & 4\mathcal{H}^{MN}\partial_M\partial_N d - \partial_M\partial_N\mathcal{H}^{MN} \\ & - 4\mathcal{H}^{MN}\partial_M d\partial_N d + 4\partial_M\mathcal{H}^{MN}\partial_N d \\ & + \frac{1}{8}\mathcal{H}^{MN}\partial_M\mathcal{H}^{KL}\partial_N\mathcal{H}_{KL} - \frac{1}{2}\mathcal{H}^{MN}\partial_M\mathcal{H}^{KL}\partial_K\mathcal{H}_{NL}\end{aligned}$$

$$S = \int dx d\tilde{x} e^{-2d} \mathcal{R}$$

Gauge Symmetry

$$\delta_\xi \mathcal{R} = \hat{\mathcal{L}}_\xi \mathcal{R} = \xi^M \partial_M \mathcal{R}$$

$$\delta_\xi e^{-2d} = \partial_M (\xi^M e^{-2d})$$

Field equations give gen. Ricci tensor

## 2-derivative action

$$S = S^{(0)}(\partial, \partial) + S^{(1)}(\partial, \tilde{\partial}) + S^{(2)}(\tilde{\partial}, \tilde{\partial})$$

Write  $S^{(0)}$  in terms of usual fields

Gives usual action (+ surface term)

$$\int dx \sqrt{-g} e^{-2\phi} \left[ R + 4(\partial\phi)^2 - \frac{1}{12} H^2 \right]$$

$$S^{(0)} = S(\mathcal{E}, d, \partial)$$

## 2-derivative action

$$S = S^{(0)}(\partial, \partial) + S^{(1)}(\partial, \tilde{\partial}) + S^{(2)}(\tilde{\partial}, \tilde{\partial})$$

Write  $S^{(0)}$  in terms of usual fields

Gives usual action (+ surface term)

$$\int dx \sqrt{-g} e^{-2\phi} \left[ R + 4(\partial\phi)^2 - \frac{1}{12} H^2 \right]$$

$$S^{(0)} = S(\mathcal{E}, d, \partial)$$

$$S^{(2)} = S(\mathcal{E}^{-1}, d, \tilde{\partial}) \quad \text{T-dual!}$$

$$S^{(1)} \quad \text{strange mixed terms}$$

- Restricted DFT:  
fields independent of half the coordinates
- If independent of  $\tilde{x}$ , equivalent to usual action
- Duality covariant: duality changes which half of coordinates theory is independent of
- Equivalent to Siegel's formulation **Hohm & Kwak**
- Good for non-geometric backgrounds

# Gauge Algebra

Parameters  $(\epsilon^i, \tilde{\epsilon}_i) \rightarrow \Sigma^M$

Gauge Algebra  $[\delta_{\Sigma_1}, \delta_{\Sigma_2}] = \delta_{[\Sigma_1, \Sigma_2]_C}$

$$[\hat{\mathcal{L}}_{\xi_1}, \hat{\mathcal{L}}_{\xi_2}] = -\hat{\mathcal{L}}_{[\xi_1, \xi_2]_C}$$

## C-Bracket:

$$[\Sigma_1, \Sigma_2]_C \equiv [\Sigma_1, \Sigma_2] - \frac{1}{2} \eta^{MN} \eta_{PQ} \Sigma_{[1}^P \partial_N \Sigma_{2]}^Q$$

Lie bracket + metric term

Parameters  $\Sigma^M(X)$  restricted to N

Decompose into vector + 1-form on N

C-bracket reduces to **Courant bracket** on N

Same covariant form of gauge algebra found in similar context by **Siegel**



# Jacobi Identities not satisfied!

$$J(\Sigma_1, \Sigma_2, \Sigma_3) \equiv [ [\Sigma_1, \Sigma_2], \Sigma_3 ] + \text{cyclic} \neq 0$$

for both C-bracket and Courant-bracket

How can bracket be realised as a symmetry algebra?

$$[ [\delta_{\Sigma_1}, \delta_{\Sigma_2}], \delta_{\Sigma_3} ] + \text{cyclic} = \delta_{J(\Sigma_1, \Sigma_2, \Sigma_3)}$$

# Symmetry is Reducible

Parameters of the form  $\Sigma^M = \eta^{MN} \partial_N \chi$   
do not act

Gauge algebra determined up to such transformations

cf 2-form gauge field  $\delta B = d\alpha$

Parameters of the form  $\alpha = d\beta$   
do not act

# Symmetry is Reducible

Parameters of the form  $\Sigma^M = \eta^{MN} \partial_N \chi$   
do not act

Gauge algebra determined up to such transformations

cf 2-form gauge field  $\delta B = d\alpha$

Parameters of the form  $\alpha = d\beta$   
do not act

Resolution:

$$J(\Sigma_1, \Sigma_2, \Sigma_3)^M = \eta^{MN} \partial_N \chi$$

$\delta_{J(\Sigma_1, \Sigma_2, \Sigma_3)}$  does not act on fields

## D-Bracket

$$[A, B]_{\text{D}} \equiv \hat{\mathcal{L}}_A B$$

$$[A, B]_{\text{D}}^M = [A, B]_{\text{C}}^M + \frac{1}{2} \partial^M (B^N A_N)$$

Not skew, but satisfies Jacobi-like identity

$$[A, [B, C]_{\text{D}}]_{\text{D}} = [[A, B]_{\text{D}}, C]_{\text{D}} + [B, [A, C]_{\text{D}}]_{\text{D}}$$

On restricting to null subspace N

C-bracket  $\rightarrow$  Courant bracket

D-bracket  $\rightarrow$  Dorfman bracket

Gen Lie Derivative  $\rightarrow$  GLD of Grana, Minasian, Petrini  
and Waldram

# Frames for Doubled Space

$$e^M_A$$

Basis, labelled by  $A=1,\dots,2D$

$$e^M_A \rightarrow e^M_B \Lambda^B_A, \quad \Lambda(X) \in GL(2D, \mathbb{R})$$

$$\mathcal{H}_{AB} \equiv e^M_A e^N_B \mathcal{H}_{MN} \quad \hat{\eta}_{AB} \equiv e^M_A e^N_B \eta_{MN}$$

e.g. Orthonormal Frame

$$e^M_A e^N_B \eta_{MN} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \rightarrow \hat{\eta}_{AB} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

Reduces tangent space group to  $O(D,D)$

# Generalized Connection Coimbra, Strickland-Constable & Waldram

$$D_M W^A = \partial_M W^A + \tilde{\Omega}_M^A{}_B W^B$$

$$\tilde{\Omega}^A{}_B W^B \in GL(2D, \mathbb{R})$$

## Generalised Lie Derivative

$$\hat{\mathcal{L}}_\xi A^N \equiv \xi^P \partial_P A^N + (\partial^N \xi_P - \partial_P \xi^N) A^P$$

## Covariantised Lie Derivative

$$\hat{\mathcal{L}}_\xi^D A^N \equiv \xi^P D_P A^N + (D^N \xi_P - D_P \xi^N) A^P$$

Difference is Covariant, defines **TORSION**

$$\hat{\mathcal{L}}_\xi^D A^N - \hat{\mathcal{L}}_\xi A^N = T_{MP}^N \xi^M A^P$$

# Generalized Curvature

$$R(U, V, W) = [D_U D_V]W - D_{[U, V]}W$$

Scale by functions:

$$U \rightarrow fU, \quad V \rightarrow gV, \quad W \rightarrow hW$$

$$R(fU, gV, hW) = fgh R(U, V, W) - \frac{1}{2}h \eta(U, V) D_{(fdg - gdf)}W$$

**Non-tensorial!**

But tensorial for vectors with  $\eta(U, V) = 0$

U, V tangent to null subspace

# The Connection

Coimbra, Strickland-Constable & Waldram

Choose conformal frames  $e^M{}_A e^N{}_B \eta_{MN} = \Phi^2 \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$

$$GL(2D, \mathbb{R}) \rightarrow O(D) \times O(D) \times \mathbb{R}^+$$

Seek torsion-Free Connection  $T^P{}_{MN} = 0$

$$D\mathcal{H} = 0, \quad D\Phi = 0$$

Find general connection **NOT UNIQUE!**

Determined up to tensor  $A$

Curvature depends on  $A$ . But  $A$  drops out of Ricci tensor, scalar curvature and Dirac equation

Field equations given by Ricci tensor, indep of  $A$

Susy variations independent of  $A$



$$\mathcal{H}^{MP} \mathcal{H}_{PN} = \delta^M_N$$

$$S^M_N \equiv \eta^{MP} \mathcal{H}_{PN} \quad \text{satisfies} \quad S^2 = 1$$

**Split basis:**

$$e^M_a \quad e^M_{\bar{a}} \quad a, \bar{a} = 1, \dots, D$$

$$S e_a = -e_a, \quad S e_{\bar{a}} = e_{\bar{a}}$$

**This form of basis preserved by**  $GL(D, \mathbb{R}) \times GL(D, \mathbb{R})$

$$\mathcal{H}_{AB} = 2 \begin{pmatrix} g_{\bar{a}\bar{b}} & 0 \\ 0 & g_{ab} \end{pmatrix}, \quad \hat{\eta}_{AB} = 2 \begin{pmatrix} g_{\bar{a}\bar{b}} & 0 \\ 0 & -g_{ab} \end{pmatrix}$$

**Partially fix gauge**

$$GL(D, \mathbb{R}) \times GL(D, \mathbb{R}) \rightarrow O(D) \times O(D) \times \mathbb{R}^+$$

$$g_{ab} = \Phi^2 \delta_{ab}, \quad g_{\bar{a}\bar{b}} = \Phi^2 \delta_{\bar{a}\bar{b}}$$

$O(D) \times O(D) \times \mathbb{R}^+$  **Torsion-Free Connection**

$$D\mathcal{H} = 0, \quad D\Phi = 0 \quad T_{MN}^P = 0$$

Take  $\Phi = e^{-2\phi} \sqrt{-g} = e^{-2d}$

**Gives non-unique connection**

$$D_a w^b = \nabla_a w^b - \frac{1}{6} H_a{}^b{}_c w^c - \frac{2}{9} (\delta_a{}^b \partial_c \phi - \eta_{ac} \partial^b \phi) w^c + A_a{}^{+b}{}_c w^c,$$

$$D_{\bar{a}} w^b = \nabla_{\bar{a}} w^b - \frac{1}{2} H_{\bar{a}}{}^b{}_c w^c,$$

$$D_a w^{\bar{b}} = \nabla_a w^{\bar{b}} + \frac{1}{2} H_a{}^{\bar{b}}{}_{\bar{c}} w^{\bar{c}},$$

$$D_{\bar{a}} w^{\bar{b}} = \nabla_{\bar{a}} w^{\bar{b}} + \frac{1}{6} H_{\bar{a}}{}^{\bar{b}}{}_{\bar{c}} w^{\bar{c}} - \frac{2}{9} (\delta_{\bar{a}}{}^{\bar{b}} \partial_{\bar{c}} \phi - \eta_{\bar{a}\bar{c}} \partial^{\bar{b}} \phi) w^{\bar{c}} + A_{\bar{a}}{}^{-\bar{b}}{}_{\bar{c}} w^{\bar{c}},$$

**Ambiguity: A terms arbitrary**

- Gualtieri:  $O(D) \times O(D)$  connection
- Waldram et al: similar  $O(D) \times O(D) \times \mathbb{R}^+$  connection, with dilaton.
- Jeon, Lee Park: similar connection, with  $A=0$
- Siegel: similar connection, but curvatures etc constructed without geometry
- Hohm & Kwak; Hohm & Zwiebach: more on Siegel construction

# Generalised Geometry, M-Theory

- Generalised Geometry doubles Tangent space, Metric + B-field, action of  $O(D,D)$  Hitchin; Gualtieri
- DFT doubles space, doubles coordinates.
- Extended geometry: extends tangent space, metric and 3-form gauge field, action of exceptional U-duality group Hull; Pacheco & Waldram
- 10-d type II sugra action in terms of extended geometry Coimbra, Strickland-Constable & Waldram Hillman
- 11-d sugra action in terms of extended geometry Berman, Perry et al Coimbra, Strickland-Constable & Waldram

# Double Field Theory

- Constructed cubic action, quartic has new stringy features
- T-duality symmetry, cocycles, symmetry a homotopy Lie algebra, constraints
- Restricted DFT: have non-linear background independent theory, duality covariant
- Courant bracket gauge algebra
- Stringy issues in simpler setting than SFT

- Full theory without restriction? Does it close on a geometric action with just these fields?
- SUSY: Jeon, Lee, Park; Hohm & Kwak; Coimbra, Strickland-Constable, Waldram,...
- Type II: Hohm, Kwak, Zwiebach; Coimbra, Strickland-Constable, Waldram; Jeon, Lee, Park, Suh,...
- Use for non-geometric backgrounds
- General spaces, not tori?
- Doubled geometry *physical* and *dynamical*

- Early work on strings and doubled space: Duff, Tseytlin...
- Sigma model for doubled geometry: Hull; Hull & Reid-Edwards,...
- Doubled sigma model: Quantization Hull; Hackett-Jones & Moutsopoulos; Beta functions: Copland; Berman, Copland & Thompson
- D-Branes: Hull; Lawrence, Schulz, Wecht; Bergshoeff & Riccioni
- Geometry with projectors; YM DFT: Jeon, Lee, Park
- $O(10,10)$  from  $E_{11}$ ; West
- Non-geometry: Andriot, Larfors, Lust, Patalong; Ditetto, Fernandez-Melgarejo, Marques & Roest
- Twisted torus: Grana & Marques; Chatzistavrakidis & Jonke
- Heterotic DFT: Hohm & Kwak
- Type II DFT: Thompson; Hull; Hohm, Kwak, Zwiebach